

## **ROTIFER DIVERSITY IN LAKE MASUNDA, THANE (MAHARASHTRA)**

Somani, Vaishali U, and Madhuri K. Pejaver,  
Zoology Department,  
B. N. Bandodkar College of Science,  
Thane - 400 601 (Maharashtra)

Key words : Rotifera, Masunda lake, Tolerant genera, Brachionus, Pollution.

### **Abstract :**

Rotifer diversity from Masunda lake, Thane (Maharashtra) was studied from April 1999 to June 2000. Rotifers constituted 34.67% of total zooplankton and their density ranged from 300 to 2000 ind/100l. The peak density was recorded in winter (November) while minima was observed in summer (May).

Dominance of family Brachionidae was observed all over the Study period, Brachionidae was observed all over the Study period, Brachionus was the dominant genera qualitatively as well as quantitatively. Dominance of this tolerant genera indicates onset of eutrophication in the ecosystem.

### **Introduction :**

Rotifera comprises an integral part in the aquatic food chain. Their role as link between the nanoplankton and carnivorous zooplankton is well established (Ruttner-Kolisko, 1974). They play a key role in cycling of organic material (Mishra and Saksena, 1998) Rotifers exhibit remarkable ability to colonize diversified freshwater biotopes and these are apparently the most sensitive indicators of water properties (Ruttner-Kolisko, 1971). Due to their short development period and fast reproductive

rate characterized by parthenogenetic production (Herzig, 198) rotifers can populate vacant niches with extreme rapidity and convert primary production into a form usable for secondary consumers, producing up to 50% of the total plankton biomass (Nogrady et al., 1993). They are very common in Indian waters and their occurrence in eutrophic water bodies is also well noted.

Rotifer population from different regions has been extensively studied by many scientists (George, 1961; Arora, 1966a; Michal, 1968; Sharma, 1992, Dhanapathi 2000). However references on rotifer diversity from fresh waters of Maharashtra,

especially in Thane-Mumbai region are scarce. Hence the study on rotifer diversity was undertaken.

### **Materials and Methods :**

The zooplankton samples were collected monthly from April 1999 to June 2000 by towing 41 µm mesh net through a boat and the samples were preserved in 4 % formalin for identification and counting. The rotifers were identified using Battish (1992) and Dhanapathi (2000).

Table 2, Monthly varifications in rotifer abundance (ind/l,) in lake Masunda, Thane, Maharashtra.

<b>1999</b>	Feb	Mar	April	May	June	Jul	Aug	Sept	Oct	Nov	
<i>B. calyciflorus</i>	0	0	62	0	90	0	1216	0	806	450	
<i>B. caudatus</i>	0	0	108	0	90	0	152	93	124	150	
<i>B. diversicomis</i>	0	0	98	121	0	34	304	651	248	150	
<i>B. falcatus</i>	0	0	0	312	1350	0	0	0	434	150	
<i>B. forficula</i>	0	0	20	20	0	0	0	0	0	200	
<i>B. angularis</i>	0	0	0	0	0	0	0	0	0	100	
<i>B. plicatilis</i>	0	0	0	0	0	0 1	0	0	0	010	
<i>keratella</i>	0	0	30	30	90	34	76	186	62	800	
<i>Testidunella</i>	0	0	0	0	0	67	76	0	0	0	
<i>Hoarella</i>	0	0	0	0	0	0	0	0	0	0	
<i>Anuropsis</i>	0	0	15	15	0	0	0	0	62	0	
<i>Asplanchna</i>	0	0	0	0	0	201	0	0	0	0	
<i>Rotatoria</i>	0	0	0	0	0	0	0	0	0	0	
<i>Epiphanes</i>	0	0	0	0	0	0	0	0	0	0	
<i>Brachionus total</i>	0	0	288	433	1530	34	1672	744	1612	1200	
<i>Rotifer total</i>	0	0	333	433	1620	336	1824	930	1736	2000	
<b>1999-2000</b>	Dec	Jan	Feb	Mar	Apr	May	Jun	Total	Aug	SD	%
<i>B. calyciflorus</i>	84	420	42	60	72	33	0	3335	2223	358.2	14.04
<i>B. caudatus</i>	210	126	84	96	108	198	216	1755	117	64.27	8.069
<i>B. diversicomis</i>	42	252	210	240	288	66	72	2776	185.1	162.9	4.246
<i>B. falcatus</i>	0	42	0	96	144	0	0	2528	168.5	352.2	6.716
<i>B. forficula</i>	0	28	84	0	0	0	72	404	26.93	55.13	6.116
<i>B. angularis</i>	0	0	0	0	0	0	72	172	11.47	30.72	0.977
<i>B. plicatilis</i>	0	56	84	40	0	33	0	213	14.2	26.53	0.416
<i>keratella</i>	126	476	252	126	108	66	0	2432	162.1	213.5	0.515
<i>Testidunella</i>	0	0	42	0	0	0	0	185	12.33	26.39	5.884
<i>Hoarella</i>	0	0	0	0	0	33	0	33	2.2	8.521	0.448
<i>Anuropsis</i>	0	0	42	15	0	33	0	167	11.13	19.43	0.08
<i>Asplanchna</i>	0	14	42	0	0	0	0	257	17.13	52.07	0.404
<i>Rotatoria</i>	0	0	42	0	0	0	0	42	2.8	10.84	0.622
<i>Epiphanes</i>	0	0	0	0	0	33	0	33	2.2	6.521	0.102
<i>Brachionus total</i>	336	924	504	532	612	330	432	11183	745.5	1050	0.08
<i>Rotifer total</i>	462	1414	924	673	720	495	432	14332	955.5	1389	34.67

**Results and Discussion :**

During the present study, total rotifer density in Masunda varied from 300 to 2000 ind/ 100 L, with minima in May 1999 and maxima during November 1999. This was comparable with total rotifer density ranging from 187.65 to 2562.8 ind / 100 L recorded by Nene (1985) in the same lake. Nene (1985) had reported rotifers as the most dominant zooplankton component of Masunda. However, in the present study, rotifers were observed to be subdominant to Copepods.

Average contribution of rotifers in total zooplankton in Masunda was (34.67%), Relatively high population densities were noticed during late monsoon and early winter noticed during the monsoon and early winter (September-November 1999) as supported by Sharma (1992). However summer maxima as reported by Khan et al. (1986) and Sharma (1992) could not be observed. Mukhopadhyay et al. (1981) and Jayadevi (1994) have reported higher rotifer densities in winter, which support rotifer peak observed in November 1999 in present study. Notable difference in quantitative abundance was seen in total rotifer population but only a single distinct peak was recorded.

Rotifera is quite a diverse group of organisms and large generic variety is observed in various lentic environments all over India. However Brachionus and Keratella are the most commonly recorded rotifer genera in Indian lakes.

During the present study, occasional presence of Epiphanes, Testudinella, Hoarella, Anuroopsis, Asplanchna and Rotaria was observed (Table 1). However, quantitatively Brachionus was the main and significantly abundant genera which was followed by Keratella in terms of

abundance and periodicity (Fig. 1). This was in accordance with observations of George (1966), Jayadevi (1994) and Hiware and Jadhav (1998). Keratella followed this in terms of abundance and regular periodicity. Predominance of one or two genera is characteristic of rotifer population (George, 1961).



**Brachionus :**

Brachionus is cosmopolitan in distribution and by far the best known genus from India. It shows large species variation and embraces 19 species with more than 14 subspecies in this country (Battish, 1992). During the present study. Brachionus was represented by 7 species as B. Brachionus was represented by 7 species as B. Calyciflorus, B. caudatus, B. divescicornis, B. falcatus, B. forficula, B. angularis and B. plicatilis, thus making it significant genera. Pejaler (1977) and Fernando (1980) mentioned Brachionus as important genera in tropical rotifer community.

Among the 7 species of this genera, Brachionus calyciflorus was the most dominant, which is

**Table 1 : Rotifers of Masunda lake.**

Order	Family	Genus and species
<i>Ploimida</i>	<i>Brachionidae</i>	<i>Brachionus calyciflorus</i>
		<i>B. divescicornis</i>
		<i>B. caudatus</i>
		<i>B. falcatus</i>
		<i>B. forficula</i>
		<i>B. angularis</i>
		<i>B. plicatilis</i>
	<i>Brachionidae</i>	<i>Keratella spp.</i>
	<i>Brachionidae</i>	<i>Anuroopsis spp.</i>
	<i>Testudinellidae</i>	<i>Testudinella spp.</i>
	<i>Trochosphaeridae</i>	<i>Hoarella spp.</i>
	<i>Asplanchnidae</i>	<i>Asplanchna spp.</i>
	<i>Epiphaniidae</i>	<i>Epiphandes spp.</i>
<i>Bdelloida</i>	<i>Philodonodae</i>	<i>Rotarja spp.</i>



supported by the observations of Patil (1978). Isairasu. (1997) and Malathi (1999). It was also reported as dominant species from take Masunde in earlier studies (Nene, 1985)

During this study, in Masunda, the density peak of *Brachionus calyciflorus* was recorded in August 1999 (1216 ind/100 L). According to Nene (1985), the spp. flourished well in the month of December and June with density of 1300.51 and 1115.13 ind/100 L respectively. Exact seasonal pattern could not be observed in abundance of *Brachionus calyciflorus* as commented by Reid and Wood (1976).

*Brachionus diversicornis* was 2nd in dominance, next to *Brachionus calyciflorus*. Though Green (1960) regarded it as mostly a summer species, Battish (1992) has reported its occurrence in monsoon (July). in northern Indian States, During the present study, this species was recorded in almost all the months of the study period and peak density was attained in the month

of september 1999 in Masunda. Earlier Nene (1985) recorded highest abundance of the species In November from the same lake.

*Brachionus falcatus* occupies 3rd position, in order of abundance in Masunda lake. The species is considered as cosmopolitan in distribution. Dhanapathi (2000) designated it as warm water species and observed it to be less variable when compared to *Brachionus calyciflorus* and *Brachionus quadridentatus*. During the present study, density of *Brachionus falcatus* varied between 0 to 1350 ind/100 L in Masunda with peak observed in June 1999. Its presence was of discontinuous nature. Malathi (1999) in Hussain sagar lake, Hyderabad, recorded this species occasionally, with quite low density (20 ind/100L) only in September.

*Brachionus caudatus* enjoyed 4th position, with average density of 117 ind/100 L. It exhibited two peaks, in December 1999 and June 2000 whereasthe density was zero in May 1999 and July 1999. Nene (1985) observed its highest abundance in January 1983 in Masunda lake. The peak value reported by Nene (1985) - 13016.4 Ind/100 L, was more than 10 times higher as compared to peak density of this species recorded in present study. Pathak and Mudgal (2002) recorded this spp. as one of the dominant rotifer member in Virla reservoir (M.P.)

*Brachionus forficula* showed lower densities in Masunda. Durve (1992) and Salaskar (1996) recorded the species, as most dominant rotifer species in Udaipur kales and Powai lake respectively. Nene (1985) however did not report its presence in the study of Masunda lake. During the present study April 1999, January, February and June 2000 were characterized by low density of this species, whereas November

1999 exhibited peak values - 200 ind/100 L in Masunda.

*Brachionus angularis* is one of the important rotifer species with respect to indication of eutrophic state. In Masunda it was recorded only twice with maximum density as 100 ind/100 L. Datta et al. (1987) noted this species in Calcutta pond and its abundance was exceptionally high, making it dominant throughout the study period. While Mudgal et al. (1990) also observed it forming sporadic blooms, Patil (1978) recorded its important contribution in winter rotifer pulse. Subbamma (1993) observed dominance of *Brachionus angularis* in temple pond study.

*Brachionus angularis* is one of the important rotifer species with respect to indication of eutrophic state. In Masunda it was recorded only twice with maximum density as 100 ind/100 L. Datta et al. (1987) noted this species in Calcutta pond and its abundance was exceptionally high, making it dominant throughout the study period. While Mudgal et al. (1990) recorded its important contribution in winter rotifer pulse. Subbamma (1993) observed dominance of *Brachionus angularis* in temple pond study.

*Brachionus plicatilis* was present only for last 4 months of study period, with highest density in February 2000-84/100 L. This species was of low density status as among other *Brachionus* species of lake Masunda. Thus genus *Brachionus* showed dominance of different species in different months, however *Brachionus calycifloris* density. Similarly the dominance of a particular spp. is not in the same every year, as dominant species in April, May and June 1999 are different from April, May and June 2000.

### **Keratella spp.**

*Keratella* was the 2nd dominant rotifer genera. This genus is also considered as cosmopolitan in distribution and according to Battish (1992) and it is probably the most common rotifer genera, as it is reported in different ecosystems all over India.

During the present study, highest density of this genus in Masunda was noted in November, in fact it was the largest contributor of total rotifer peak of this month. Jayadevi (1994) and Malathi (1999) recorded *Keratella tropica* as common eurytopic perennial form in lakes of Hyderabad. In Dhom reservoir, Maharashtra (Trivedi, 1993) *Keratella* was recorded as most dominant rotifer genera. However, it was recorded only occasionally in Salim Ali lake, Aurangabad (Hiware and Jadhav, 1998)

*Anuroopsis*, the important indicator species (Gannon and Stemberger, 1978) was represented in 5 months in Masunda with low densities.

### **Other rotifers :**

The other rotifer genera (Table 2) were noted occasionally with low densities, except *Asplanchna*. High density of this genus was noted in July 1999 (201 Ind / 100 L.)

### **Egg bearing Rotifers :**

Rotifers are mostly parthenogenic. As rotifers are characterized by fast reproduction, the number of egg bearing rotifer individuals during certain period may reflect the reproductive behaviour of rotifers.

During the present study, the egg bearing rotiferan females were counted as gravid females. The counting was done in

Brachionus and Keratella, the dominant rotifers observed. (Fig. 2)

Different species of Brachionus showed the presence of gravid females, Brachionus diversicornis was observed more frequently in all seasons i.e. summer, monsoon and winter, with density varying from 4 - 186 (Ind/L. (Fig. 3) While gravid females of other species were seen only in few months as Brachionus calyciflorus in October 1999 and January 2000, with number varying from 62-228 ind/100L. Brachionus caudatus in winter, Brachionus falcatus in June, October and January, Brachionus plicatilis showed the lowest density of gravid females as 42 ind/100L. While Brachionus fucula and Brachionus angularis never showed the presence of gravid females. Gravid females of Keratella spp. in Masunda were specific to winter. Nene (1985) noted, total rotifer maxima in the same month as well as total berried rotifer maxima (April), which was not observed during the present study.

Brachionus calyciflorus and Brachionus diversicornis the dominant species in the lake, showed more number of their gravid individuals. Winter 2000 was characterized by presence of gravid females of 6 rotiferan types, Brachionus calyciflorus, B. caudatus, B. diversicornis, B. falcatus, B. plicatilis and Keratella spp. Similarly the peak density of rotifers was also noted in winter. Hence winter seems to be favorable for reproduction of rotifers in Masunda.

The present study also suggests that genus Brachionus is a continuous breeder and hence one or the other species of Brachionus breeds in every month resulting in the dominance of this genera.

#### **Acknowledgement :**

The authors wish to thank the Principal and staff of zoology department, B. N. Bandodkar College of Science, for their constant encouragement.

#### **References :**

Arora H. C. (1966 a) - Rotifers as indicators of trophic nature of environment, Hydrobiologia 27, pp : 146 -159

Battish, S. K. (1992) Freshwater zooplankton of India Oxford & IBH Publishing Co. pp : 233

Datta, N. C., Mandal, N. and Bandopadhyay (1987) Seasonal abundance of rotifers in a perennial freshwater pond in Calcutta J. Environ. Biol 8(I) pp : 63 - 71

Dhanapathi, M.V.S.S.S. (2000) Taxonomic notes on the Rotifers from India - IAAB Publication, Hyderabad pp : 175

Durve (1993) Past, present and future of Udaipur lakes as a source of drinking water. In - Recent Advances in Freshwater Biology

Fernando, C. H. (1980) The freshwater zooplankton of Srilanka, with a discussion of tropical freshwater zooplankton composition. Int. Revue ges Hydrobiol, 45 - pp : 85 - 125

Gannon, John E. and R. S. Stemberger (1978) Zooplankton (especially crustaceans and rotifers) as indicators of water quality. Trans. Amer. Micros. Soc. Vol. 97 No. I Jan pp : 16-32

George, M. G. (1961) Observation on rotifer from shallow ponds in Delhi Curr Sci. (30) pp : 247 : 250

Green, j. (1960) Zooplankton of river Skoto. The Rotifera Proc. Zool. Soc. London, 135, pp : 491 - 523

Herzig, A. (1983) Comparative studies on the relationship between temperature & duration of embryonic development of rotifers. Hydrobiologia. 104pp - 237 - 246

Hiware, C. J. and B. V. Jadhav (1998) Studies on the microzootic fauna of Salim All Lake at Aurangabad, Maharashtra, India.

J. Aqua. Biol. Vol. 13 ( 1 & 2) pp : 18 - 21

Isalarasu, L. (1997) A report on the most common zooplankton observed in three ponds within ANJAC Campus - ANJAC Journal 14, 41 (1997)

Jaya Devi M. (1994) - Seasonal variation and population density of rotifers in three trakes of Hyderabad. : Journal of Aquatic Biology volume 9 (1 & 2) pp : 41 -44

Malathi, D. (1999) Ecological studies on lake Hussain Sagar with special referance to the zooplankton communities - Ph. D. Thesis Osmania University, Hyderabad.

Michael, R. G. (1968) Studies on the zooplankton of a tropical fish pond, Hydrobiologia 23, pp - 47- 48

Mishra, S. R. and D. N. Saksena (1998) Rotifers and their seasonal variation in a sewage collecting Morar (Kalpi) river, Gwalir, India J. Environ., Biol. 19 (4) pp - 363-374

Mudgal, H. (1990) Limnological status of a sewage polluted pond in an industrial township-In-Recent trends in Limnology Name of Journal, Pages ?

Mukhopadhyay, S. K., Babu Rao M., Maley S. V. and Yadav B. E. (1981) A Study of the rotiferan population from Waghhol, Poona, Proc. Symp. Ecol. Anim. Popul. Zool. Surv. India, Pt 2 pp : 47 - 62

Nene Vrinda N. (1985) Ecological studies of Masunda lake Thane. M.Sc. Thesis submitted to University of Mumbai (unpublished) pp -156

Nogrady, T., Wallace, R. L. & Snell, T. W. (eds) (1993) - Rotifera - guides to the identification of the microinvertebrates of he continental waters of the world, 4 SPB Academic Publishing The Hague, pp : 142

Pathak, S. K. and L. K. Mudgal (2002) A preliminary survey of Virla reservoir of Khargone (Madhy Pradesh) India, Indian Journal of Environment and Ecoplanning Vol. 6 (2) pp : 297 - 300

Patil, S. G. (1978) Plankton ecology of a few water bodies from Nagpur. Ph. D. Thesis submitted to University of Nagpur.

Pejler, B. (1977) On the global distribution of the family Brachionidae (Rotatoria). Arch. Hydrobiol. Suppl., 53 pp : 255-306

Reid G.K, and wood R. D. (1976) Ecology of inland Thresh, J. C., Sucking, E.V. & Beale, J. e. 1944 - The examination of water and water supplies, London.

Ruttner - Koliskn, A. (1971) Rotatorien also indikatoreu Furden Chemistrus Von Binnesalzgewassern Sitz, Ber. Osteff. Akad. Wiss. Math. Nat. KI. Abt. 1, 179 pp L 283 - 298

Ruttner-Koliskn, A. (1971) Rotatorien also indikatoreu Furden Chemistrus Von Binnesalzgewassern sitz. Ber. Osteff. Akad. Wiss. Math. Nat. KI. Abt. 1, 179 pp : 283 - 298

Ruttner-Koliskn, A. (1974) Planktonic rotifers Biology and Taxonomy. Schw. Verlag. Stuttgart pp : 146

Salaskar Pramod (1996) Environmental Studies of Powal Lake. Ph. D. Theissi submitted to University of Bombay.

Sharma, B. K. (1992) Systematics, Distribution & Ecology of Freshwater

rotifera in West Bengal - Aquatic Ecology  
Ashish Publishing House, New Delbl.

Subbamma, D. V. (1993) Studies on a  
plankton of a fish pond in Kolleru Lake  
area, Andhra Pradesh, India. J. Aquatic  
Biology Vol. 8(1&2) pp : 10 - 13)

Trivedy, R. K. (1993) Water quality of Dhom  
reservoir MAharashtra, India - Ecology and  
pollution of indian Lakes and Reservoirs  
(Ed) P. C. Mishra and R. K. Trivedy, Ashish  
Publishing House pp : I - 25.